

An Instance of Tick Feeding to Repletion Inside a Human Nostril

Gary P. Aronsen¹ and Richard G. Robbins²

¹ Department of Anthropology, Yale University, P. O. Box 208277, New Haven CT 06520-8277 USA
— email: gary.aronson@yale.edu

² DPMIAC / AFPMB, Walter Reed Army Medical Center, Washington DC 20307-5001 USA
— email: richard.robbs@osd.mil

ABSTRACT

A single fully engorged nymphal tick of the genus *Amblyomma* was collected from the nostril of a field researcher returning from Kibale National Park, Uganda. While unfed ticks have previously been removed from human nostrils, this is the first time that a specimen so situated was permitted to feed to repletion and detach naturally.

KEYWORDS

Tick, *Amblyomma*, human nostril, Uganda.

Introduction

Ticks (Acari: Ixodoidea) are a relatively small group (about 850 species worldwide) of large (adults 2 to 30 mm in length), obligately hematophagous, ectoparasitic mites of paramount importance in the transmission of disease among domestic and wild mammals; they are second only to mosquitoes as vectors of human disease (Sonenshine et al. 2002). Humans, by virtue of their sophisticated learning and grooming abilities, seldom suffer prolonged tick attachment on parts of the body that are subject to regular visual examination or palpation; among those that do, many are inexperienced children. Thus, cases of tick parasitism of human eyelids, or the eyeball itself, are infrequent (Miyamoto and Miyamoto 1990; Singh et al. 2006), while cases of otocariasis — parasitism of the ear canal — are more often seen by veterinarians than by physicians

(Keirans and Pound 2003). We are aware of just a single record of an attached, blood-engorged tick being mistaken for a hemorrhoid (Chang 1958a, 1958b; see notes on this specimen in Robbins and Robbins 2003). Here we report what we believe to be only the second tick collection from human nostrils, and the first in which a tick so situated was able to feed to repletion.

In 2006, a team from the Yale University Department of Anthropology began long-term research on primates and elephant ecology at Mainaro (lat 0°21'35"N, long 30°23'18"E), a site in Uganda's Kibale National Park, one of the largest (766 km²) protected forests in Uganda. This park is home to multiple species listed as endangered or vulnerable by the International Union for Conservation of Nature and Natural Resources and the Convention on International Trade in Endangered Species of Wild Fauna and Flora (Struhsaker 1997; IUCN 2007). The

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE OCT 2008		2. REPORT TYPE		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE An Instance of Tick Feeding to Repletion Inside a Human Nostril				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Defense Pest Management Information Analysis Center (DPMIAC), Armed Forces Pest Management Board (AFPMB), Walter Reed Army Medical Center, Washington, DC, 20307-5001				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			




FIGURE 1. Nymphal *Amblyomma* (YPM 412973). A. Dorsal view. B. Ventral view. C. Capitulum and scutum. D. Legs (note coxal spurs).

Mainaro site consists of undulating hills and valleys (elevation about 1200 m) covered in a mosaic of vegetation types, including unlogged mature forest, former homesteads undergoing natural succession, and several areas where succession has been arrested because of elephant activity (Struhsaker et al. 1996; Chapman et al. 1997; Struhsaker 2003; Aronsen and Teelen 2007).

Tick infestation occurred during June 2008 fieldwork at the Mainaro site, while Aronsen was conducting vegetative surveys, primate censuses, chimpanzee nest counts and group follows of gray-cheeked mangabeys (*Lophocebus albigena* (Gray)) in both mature forests and disturbed or reforested areas. On about 29 June 2008, Aronsen noted an unusual sensation in his right nostril and determined that a tick had attached itself

there. Because he was about to depart Uganda, he decided not to extract the tick until his return to the United States, but on 4 July 2008, during a stopover in Amsterdam, the Netherlands, a fully engorged tick was expelled into his hand by a strong sneeze. The specimen was retained and deposited on 24 July 2008 in the Division of Entomology of the Peabody Museum of Natural History, Yale University (catalog number YPM 412973). Subsequently, it was shipped to the Walter Reed Army Medical Center, Washington, DC, USA, where Robbins identified it as a nymph of the pantropical genus *Amblyomma*. On its return to the Yale Peabody Museum, the tick's diagnostic morphological characters were photographed (Figure 1).

Ticks commonly infest the nostrils or nasal cavities of reptiles, birds and mammals, irritating

or incapacitating their hosts and transmitting a range of infectious agents (Amerson 1966; Gladney et al. 1972; Hesse 1985; Cunha et al. 2007). Apparently the only other record of ticks collected from the nostrils of humans is that of Walton (1960), who encountered this phenomenon in the Bunyoro and Kayonza ("Impenetrable") Forests, also in Uganda. Like our case, Walton's tick specimens were nymphal *Amblyomma*, but were described as white, meaning that they had not begun to imbibe blood, although their human hosts reported a mild tenderness when the nose was touched or blown. Walton (1960) speculated that this insignificant reaction might occur if humans were serving as alternate hosts for a tick species that usually infests an anthropoid ape, such as the chimpanzee (*Pan troglodytes* (Blumenbach)). However, while *Amblyomma cohaerens* Dönitz has occasionally been collected from Ugandan *P. troglodytes* and the hamadryas baboon (*Papio hamadryas* (Linnaeus)), this tick chiefly parasitizes the African buffalo (*Syncerus caffer* (Sparrman)) and warthog (*Phacochoerus africanus* (Gmelin)) (Matthysse and Colbo 1987). The former species is common within Kibale. Moreover, subsequent collecting has shown that African primates are more often hosts of the unrelated tick genus *Rhipicephalus*, 12 species of which have been recorded from primates in Africa — exactly one-fifth of the African *Rhipicephalus* fauna (Walker et al. 2000). Because the preimagines (larvae, nymphs) of most Afrotropical *Amblyomma* cannot be determined to species using morphological characters (Robbins, unpublished data), the species responsible for human nasal parasitism in Uganda will remain unknown pending formal rearing studies or molecular genetic analysis.

Acknowledgments

We thank the Uganda Wildlife Authority, the President's Office, and the Uganda Council of Science and Technology for permission to conduct research in Kibale National Park. Thanks also to our co-principal investigators, David Watts and Simone Teelen, and to Assimwe Dan and Birungi Patrick for field assistance. We are especially grateful to Eric Lazo-Wasem and Lawrence F. Gall for taking the photographs and for editorial advice, and Leonard E. Munster-

mann, Raymond J. Pupedis and William L. Krinsky for their interest in this tick specimen, as well as their helpful counsel and logistical support.

Received 30 August 2008; revised and accepted 4 September 2008.

Literature Cited

- AMERSON, A. B. 1966. *Ornithodoros capensis* (Acarina: Argasidae) infesting sooty tern (*Sterna fuscata*) nasal cavities. *Journal of Parasitology* 52:1220-1221.
- ARONSEN, G. P. AND S. TELEN. 2007. New research at Mainaro, Kibale Forest National Park, Uganda: primate and other mammal diversity in a mixed regenerating forest. *American Journal of Physical Anthropology* S42:65.
- CHANG, P. H. 1958a. A study of *Amblyomma cyprum* [actually *A. testudinarium* Koch] collected from human body. *Acta Entomologica Sinica* 8:290-292. Chinese.
- 1958b. A case of tick infestation in the anus of the human body. *National Medical Journal of China* 44:304-305 (in Chinese).
- CHAPMAN, C. A., L. J. CHAPMAN, R. WRANGHAM, G. ISABIRYE-BASUTA AND K. BEN-DAVID. 1997. Spatial and temporal variability in the structure of a tropical forest. *African Journal of Ecology* 35:287-302.
- CUNHA, A. P. DA, A. C. P. DE P. BELLO, R. C. LEITE, A. C. C. L. RIBEIRO, C. M. V. DE FREITAS, E. BASTIANETTO AND P. R. OLIVEIRA. 2007. Efeito do controle estratégico de *Amblyomma cajennense* (Fabricius, 1787) (Acari: Ixodidae) sobre a população de *Anocentor nitens* (Neumann, 1897) (Acari: Ixodidae) em equinos. *Revista Brasileira de Parasitologia Veterinária* 16:215-219.
- GLADNEY, W. J., S. E. ERNST, C. C. DAWKINS, R. O. DRUMMOND AND O. H. GRAHAM. 1972. Feeding systemic insecticides to cattle for control of the tropical horse tick. *Journal of Medical Entomology* 9:439-442.
- HESSE, G. H. 1985. Interstadial competition for sites of attachment to hosts in a one-host reptile tick in Senegal. *Acarologia* 26:355-359.
- [IUCN] INTERNATIONAL UNION FOR CONSERVATION OF NATURE AND NATURAL RESOURCES. 2007. 2007 IUCN Red List of Threatened Species [internet]. Cambridge, UK: IUCN, SSC Red List Programme; [accessed 2008 August 25]. Available from: <http://www.iucnredlist.org/>
- KEIRANS, J. E. AND J. M. POUND. 2003. An annotated bibliography of the spinose ear tick, *Otobius megnini* (Dugès, 1883) (Acari: Ixodida: Argasidae) 1883-2000. *Systematic and Applied Acarology Special Publications* 13:1-68.
- MATTHYSSE, J. G. AND M. H. COLBO. 1987. The Ixodid Ticks of Uganda, Together with Species Pertinent to Uganda Because of Their Present Known Distribution. College Park, MD: Entomological Society of America. 426 pp.
- MIYAMOTO, K. AND K. MIYAMOTO. 1990. The first case of human eyeball infestation with an ixodid tick in Japan. *Japanese Journal of Sanitary Zoology* 41:273-274.
- ROBBINS, R. G. AND E. M. ROBBINS. 2003. An indexed, annotated bibliography of the Chinese- and Japanese-language papers on ticks and tick-borne diseases translated

- under the editorship of the late Harry Hoogstraal (1917–1986). Systematic and Applied Acarology Special Publications 17:1–12.
- SINGH, M., P. GOPALAKRISHNAKONE AND R. L. S. YEOH. 2006. Ixodes tick infestation of the eyelid of a child. Canadian Journal of Ophthalmology 41:783–784.
- SONENSHINE, D. E., R. S. LANE AND W. L. NICHOLSON. 2002. Ticks (Ixodida). In: G. Mullen and L. Durden, eds. Medical and Veterinary Entomology. New York: Academic Press. pp. 517–558.
- STRUHSAKER, T. T. 1997. Ecology of an African Rain Forest: Logging in Kibale and the Conflict between Conservation and Exploitation. Gainesville, FL: University Press of Florida. 432 pp.
- 2003. Evaluation of the UWA–FACE Natural High Forest Rehabilitation Project in Kibale National Park, Uganda. Arlington, VA: CABS. 15 pp. Unpublished report prepared for the Center for Applied Biodiversity Science of Conservation International and for the FACE Foundation. Available from the CABS.
- STRUHSAKER, T. T., J. S. LWANGA AND J. M. KASENENE. 1996. Elephants, selective logging and forest regeneration in the Kibale Forest, Uganda. Journal of Tropical Ecology 12:45–64.
- WALKER, J. B., J. E. KEIRANS AND I. G. HORAK. 2000. The Genus *Rhipicephalus*: A Guide to the Brown Ticks of the World. Cambridge, UK: Cambridge University Press. 643 pp.
- WALTON, G. A. 1960. A tick infesting the nostrils of man. Nature 188:1131–1132.